

PATIENT SELECTION TOOL

CHEETAH NICOM & STARLING SV

PATIENT SELECTION TOOL

Patient Type

- Shock States/Low Blood Pressure: Sepsis, Low Vascular Tone, Low Cardiac Output, Hypovolemia, Neurogenic Shock¹
- Patients treated with Inotropes, Vasopressors or Vasodilators.¹
- Surgical Patients: Perioperative Volume Management, Goal Directed Therapy, Enhanced Recovery After Surgery (ERAS)²
- Emergency/Trauma Patients³
- Other Critical Care Conditions: Acute Respiratory Distress (ARDS),⁴ Sub-Arachnoid Hemorrhage (SAH),⁵ Acute Kidney Injury (AKI),⁶ and Congestive Heart Failure (CHF)⁷
- Patients undergoing Continuous Renal Replacement Therapy (CRRT) or patients undergoing hemodialysis⁸

ONLY - 50% OF HEMODYNAMICALLY UNSTABLE PATIENTS WILL RESPOND TO FLUID BY INCREASING CARDIAC OUTPUT AND PERFUSION.⁹

References: 1. Marik P et al. The use of bioreactance and carotid Doppler to determine volume responsiveness and blood flow redistribution following passive leg raising in hemodynamically unstable patients. *Chest* 2013; 143:364-370. 2. Waldron N et al. A prospective comparison of a noninvasive cardiac output monitor versus esophageal Doppler monitor for goal directed fluid therapy in colorectal surgery patients. *Anesth Analg* 2014; 118:966-75. 3. Dunham, CM et al. Emergency department noninvasive (NICOM) cardiac outputs are associated with trauma activation, patient injury severity and host conditions and mortality. *J Trauma Acute Care Surg*. 2012; 73:479-85. 4. National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network. Comparison of two fluid management strategies in acute lung injury. *New Engl J Med* 2006; 254:2564-2575. 5. Mittal M et al. Management of catecholamine-induced stunned myocardium — a case report. *J Clin Anesth* 2015; 27:527-30. 6. Grams ME et al. Fluid balance, diuretic use, and mortality in acute kidney injury. *Clin J Am Soc Nephrol* 2011; 6:966-973. 7. Maurer MM et al. A multicenter study of noninvasive cardiac output by bioreactance during symptom-limited exercise. *J Card Fail* 2009; 15:689-99. 8. Kossari N et al. Bioreactance: a new tool for cardiac output and thoracic fluid content monitoring during hemodialysis. *Hemodial Int* 2009; 13:512-7. 9. Michard F and Teboul JL. Predicting Fluid Responsiveness in ICU patients: a critical analysis of the evidence. *Chest*. 2002; 121:2000-2008.



Cheetah Medical, Inc.

600 SE Maritime Ave Suite 220, Vancouver, WA 98661 USA
Toll free: 866.751.9097 • Tel: 360.828-8685 • cheetah-medical.com

Cheetah NICOM and Starling SV are trademarks of Cheetah Medical, Inc.
©2016 Cheetah Medical

R-MRK-060